

Systematic Review

The Top 50 Most Cited Articles on Iliotibial Band Syndrome: a bibliometric analysis

Abdullah B Chandasir, BS^{1a}, Noora S Chandasir, BA^{2b}, Yagni Patel, BS^{1c}, Akshar Patel, BS^{1d}, Umar Ghilzai, MD^{3e}, Phillip N Williams, MD^{3f}

¹ Medical College of Georgia, ² Philadelphia College of Osteopathic Medicine, ³ Department of Orthopaedic Surgery, Baylor College of Medicine

Keywords: Iliotibial Band Syndrome, Citation Analysis, Sports Medicine, Bibliometric Analysis

<https://doi.org/10.60118/001c.145737>

Journal of Orthopaedic Experience & Innovation

Vol. 7, Issue 1, 2026

Background

Iliotibial band syndrome (ITBS) is a common overuse injury that is found predominantly in athletes that undergo extensive repetitive lower extremity movement such as runners and cyclists. Despite extensive literature relating to ITBS, the quality of evidence and research trends remain unclear.

Methods

A literature search was conducted using the Web of Science platform to identify the top cited papers on IT band syndrome. The final selection of 50 English-language articles was

a Abdullah Chandasir is a current 3rd year at the Medical College of Georgia. He is interested in orthopedic surgery, and is involved in research, mentorship and advocacy in orthopaedics.

[Conflicts of Interest Statement for Abudallah Chandasir](#)

b Noora Chandasir is a medical student whose prior education and research centered on human rights, with a focus on torture and ill-treatment of political prisoners. Her background includes work involving organ trafficking and nuclear disarmament, as well as field-based human rights research with refugee communities through HRMI, and alongside partners such as Human Rights Watch. She hopes to build a career that draws on both medicine and her longstanding commitment to humanitarian and global issues

[Connect with Noora Chandasir on LinkedIn](#)

[Conflicts of Interest Statement for Noora Chandasir](#)

c Yagni Patel is a 4th year medical student at the Medical College of Georgia, and went to Emory University for undergraduate school.

[Conflicts of Interest Statement for Yagni Patel](#)

d Akshar Patel is a third-year medical student at the Medical College of Georgia. He is from Toccoa, Georgia, and will be going into anesthesiology.

[Conflicts of Interest Statement for Akshar Patel](#)

e Dr. Ghilzai is a fourth year orthopaedic surgery resident at the Baylor College of Medicine, and is interested in hand and upper extremity surgery.

[Conflicts of Interest Statement for Dr. Umar Ghilzai](#)

[Visit the Open Payments Data Page for Dr. Umar Ghilzai](#)

f Phillip Williams, MD is a board-certified orthopedic surgeon specializing in sports medicine in the Joseph Barnhart Department of Orthopedic Surgery at Baylor Medicine. Dr. Williams earned an undergraduate degree in Political Science at Yale while competing on the track & field team. Afterward, he attended Harvard Medical School, followed by an orthopedic surgery residency at Hospital for Surgery in New York. Dr. Williams completed additional training in Sports Medicine that included a fellowship at Cedars-Sinai Kerlan-Jobe Institute, where he had the opportunity to assist in the care of several professional sports teams, including the Los Angeles Lakers, Los Angeles Dodgers, Los Angeles Kings, Anaheim Ducks, Los Angeles Sparks, and the Los Angeles Galaxy.

His clinical and research interests include ACL injury and prevention, joint preservation, biologics, and rotator cuff conditions. He utilizes novel techniques and innovations to relieve pain, restore function, and confidently return patients to their desired activities.

[Visit Dr. Phillip Williams](#)

[Connect with Dr. Phillip Williams on LinkedIn](#)

[Conflicts of Interest Statement for Dr. Phillip Williams](#)

[Visit the Open Payments Data Page for Dr. Phillip Williams](#)

analyzed for journal, author location, citation count, publication year, citation density, general topic, and level of evidence (LOE). Kruskal-Wallis tests assessed citation differences across categories, followed by Dunn's tests when appropriate ($p < 0.05$).

Results

The top 50 most cited papers on iliotibial band syndrome (ITBS) had a total of 3,938 citations (mean 78.8 ± 75.5). The average citation density was 4.3 ± 3.7 , with a maximum of 18.4. Publication years ranged from 1980 to 2018, with 28% published before 2000. These papers appeared in 32 journals, most commonly in The American Journal of Sports Medicine (12%). Authors from the United States contributed to 62% of the publications. The most common levels of evidence were IV (46%) and V (38%). Dominant topics included anatomy/pathophysiology (32%) and biomechanics/gait analysis (32%). Citation density differed significantly by topic ($p = 0.033$), with biomechanics/gait analysis papers receiving more citations than those on diagnosis/imaging ($p = 0.032$) and surgical management ($p = 0.011$). Citation density was also positively associated with the more recent publication year ($p = 0.04$).

Conclusion

The most influential literature on ITBS is largely composed of older, lower-level evidence studies, with a predominant focus on anatomy and biomechanics. Despite advances in sports medicine, high-level evidence on ITBS remains limited. These findings underscore the need for more contemporary, high-quality studies to guide evidence-based management of ITBS.

INTRODUCTION

Iliotibial band syndrome (ITBS) is a common overuse injury that is found predominantly in athletes that undergo extensive repetitive lower extremity movement such as runners and cyclists. ITBS results from friction or compression of the lateral iliotibial band against the lateral femoral epicondyle, leading to inflammation and lateral knee pain, and is influenced by biomechanical factors such as weakness or inhibition of the lateral gluteal muscles, with additional risk factors including excessive running or cycling and sudden changes in running surfaces (Pegrum, Self, and Hall 2019). Recent studies have associated female sex, younger age, fewer years of running experience, and faster running speed with ITBS (Marais et al. 2024). ITBS primarily affects younger and middle aged patients, with the average age of patients being 31 years (Noble 1980). Recent studies have found IT band syndrome to be present in 5% - 14% of runners, highlighting its significance as a common athletic ailment. Treatment initially involves conservative methods, consisting of rest, stretching, pain management, and modification of running habits (Beals and Flanigan 2013). Corticosteroid injections can be utilized as a second line treatment modality for persistent pain, and surgery is rarely performed in refractory cases (Beals and Flanigan 2013). The incidence and management of ITBS can result in significant stress in patients, especially those looking to return to high-level activity or employment. Thus, new treatment modalities and guidelines are paramount in the return of patients to activity.

One way to monitor the direction and trends of new research in IT band syndrome, and medicine in general, is bibliometric analyses. Bibliometric analyses allow for the identification of recent developments and emerging trends within a certain field, as well as offering insights into po-

tential directions for future investigation (Ellegaard 2018). They also identify the leading researchers and institutions in the field of interest, allowing for potential collaboration and the sharing of ideas (Ellegaard 2018).

These studies identify and sort the top papers by citation and citation index, or citations in relation to publication year. This type of analysis facilitates the identification of the most influential publications within the field, along with their defining characteristics. Although there has been a plethora of bibliometric analyses performed on the most common orthopedic ailments, there has not been a bibliometric analysis done on the literature on ITBS.

This study aims to identify the 50 most-cited publications on iliotibial band syndrome and to evaluate their key characteristics and thematic focus. We predict that the bulk of these publications will be older than 10 years in age and will predominantly represent lower levels of evidence.

METHODS

A literature search was carried out in January 2025 using the Web of Science Core Collection platform (Clarivate Plc, Boston, MA, USA). The search terms "iliotibial band syndrome" or "IT band syndrome" were utilized, and the search was queried and sorted in descending order by citations. Two of the authors then analyzed the titles and abstracts of the preliminary search, and sorted them based on specific inclusion/exclusion criteria. Articles were excluded if they were unrelated to IT band syndrome or if they were not in English. Articles were included if they were pertaining to anatomy/pathophysiology of ITBS, risk factors and epidemiology, diagnosis and imaging modalities for ITBS, biomechanics and gait analyses of patients with ITBS, treatment and rehabilitation, surgical management, strength and conditioning, or case reports of ITBS.




Figure 1. PRISMA Flow Diagram of Study Selection

Exclusion and inclusion were performed individually, then lists were compared (Figure 1). Any discrepancies were discussed, and a third author was consulted if needed. The finalized list of 50 publications was then exported to Microsoft Excel (Microsoft Corporation, Redmond, WA). Each paper was then assigned a level of evidence according to the previously established criteria in JBJS by Wright et. Al (Wright, Swiontkowski, and Heckman 2003). For each publication, the journal, geographic location of the authors, number of citations, publication year, citation density (number of citations divided by number of years since publication), general topic of the publication, and level of evidence (LOE) were recorded. The Excel sheet was then imported into R studio version 2024.12.1-563 for statistical analysis. Statistical analysis was performed using Kruskal-Wallis tests to determine if there were any significant differences in citations between categorical variables. Dunn's test was done if Kruskal-Wallis test p value was < 0.05 to delineate specific differences among groups.

RESULTS

The top 50 cited papers had a total number of citations of 3938, and a mean number of citations of 78.8 citations per

paper (SD = 75.5). The range of citations among the top 50 most cited papers was broad; the most cited paper from our search had 461 citations, whereas the least cited paper had 22 citations. The average citation density was 4.3 (SD = 3.7). The greatest citation density was 18.4 and belonged to a paper by Fredericson et. al (Fredericson et al. 2000). The publication years ranged from 1980 to 2018; 14 (28%) of the papers were published before 2000. The years 2007, 2008, 2011, and 2014 all had 4 (8%) publications among the top 50 most cited papers, which was the most for a single year (Figure 2). The top 50 most cited papers were published across 32 journals. The American Journal of Sports Medicine (AJSM) had the most publications among these journals, with 6 (12%) of the included studies being published in AJSM, followed by The Scandinavian Journal of Medicine and Science in Sports with 4 (8%) studies. The majority of papers were published by authors based in the United States (31;62%) followed by the United Kingdom (n = 4;8%). Dr. Michael Fredericson from Stanford Orthopaedic Surgery was involved in the authorship of the greatest number of included papers, with 7 papers. Among the included papers, the most common LOE was level IV (n = 23;46%) followed by level V (n = 19;38%) (Figure 3). Of the general topics covered by the articles, the most common topics were anatomy




Figure 2. Distribution of the Top 50 Most Cited Publications on IT Band Syndrome by Publication Year

and pathophysiology of ITBS ($n = 16$; 32%) and biomechanics and gait analysis ($n = 16$; 32%) (Figure 4).

When looking at the variables that could affect the number of times these papers were cited, there was a statistically significant difference in citation density found among the different general topics of papers ($p = 0.033$). Papers about biomechanics and gait analysis were found to have more citations than papers about diagnosis and imaging ($p = 0.032$) and surgical management ($p = 0.011$). Papers about biomechanics and gait analysis also trended towards more citations than case reports ($p = 0.061$) and papers on treatment and rehabilitation ($p = 0.07$), however neither of these findings were statistically significant (Figure 5). There was also a correlation found between publication year and citation density; papers that were published more recently were found to have a greater citation density ($p = 0.04$) (Figure 6).

There was no statistically significant difference between total number of citations and general topic ($p = 0.42$), country of origin ($p = 0.73$), level of evidence ($p = 0.67$), publication year (0.60), or journal ($p = 0.30$). There were also no statistically significant differences in citation density and country ($p = 0.75$), LOE ($p = 0.21$), and journal ($p = 0.21$).

DISCUSSION

Our analysis on the top 50 most cited papers on Iliotibial band syndrome confirmed that most of the most cited papers on IT band syndrome are relatively recent, with many being published in the last 15 years. This study also found a relative lack of articles with higher LOE among the most cited articles. Furthermore, our study indicated a pronounced emphasis on topics such as biomechanics and gait




Figure 3. Distribution of the top 50 Most Cited Publications on IT Band Syndrome by Level of Evidence

analysis, as well as anatomy and pathophysiology. Papers on biomechanics/gait analysis also garnered a significantly higher number of citations compared to those addressing other aspects of ITBS.

The most cited paper on ITBS was a case series by Fredericson et. al published in 2000, with a total citation number of 461 and a citation index of 18.4 (Fredericson et al. 2000). This study aimed to measure hip abductor strength in long-distance runners with ITBS in both affected and unaffected legs, and compare strengths of both limbs, as well as to limbs of healthy long-distance runners. This study found that hip adductor strength in affected limbs was significantly weaker than strength in unaffected limbs; strengths in both limbs was significantly lower than that in control healthy patients. Fredericson et. al also found that most of the subjects with ITBS became pain free after undergoing physical therapy designed to strengthen hip abductor muscles. This study was groundbreaking as it provided early evidence linking hip abductor weakness to ITBS and suggesting targeted therapy. Its findings may have been a source of its high citations and citation index, as other authors sought to expand and discuss these novel findings.

Among the papers included in our study, the most prevalent topic categories were “biomechanics and gait analysis” and “anatomy and pathophysiology”. The “biomechanics and gait analysis” category was also found to have significantly more citations than many other categories. The most cited papers in the “biomechanics and gait analysis” category were a prospective cohort study by Noehren et. Al (Noehren, Davis, and Hamill 2007) conducting a gait analysis in healthy female recreational runners and a cross-sectional study analyzing hip, knee, and ankle 3D kinemat-




Figure 4. Distribution of the Top 50 Most Cited Publications on IT band Syndrome by General Topic




Figure 5. Citation Density by General Topic

ics in female patients by Ferber et. al (Ferber et al. 2010). The most cited papers in the “anatomy and pathophysiology” category were the previously mentioned study by Fredericson et. al (Fredericson et al. 2000) and a cadaveric study performed by Fairclough et. al (Fairclough, Hayashi, Toumi, et al. 2006). As early hypotheses on the cause of ITBS around abnormal lower-limb mechanics, developing capabilities in gait analysis allowed for the establishment of confirmatory links between running mechanics and even-




Figure 6. Correlation between Publication Year and Citation Density

tual pathology. Noehren et. al was among the first to establish the fact that the improper mechanics found in females with ITBS were present before the onset of symptoms, thus establishing temporality, and proposing that the improper mechanisms of movement in patients with ITBS were not due to the pain, but rather a possible cause of the pathology itself (Noehren, Davis, and Hamill 2007). High citation counts of this study as well as others in this cate-

gory point to the importance of biomechanical studies in determining treatments and causes of pathologies. Equally important were studies that clarified the pathophysiology of ITBS, particularly by reshaping perceptions of its underlying causes. Fairclough et al. in particular established through MR-based analysis as well as gross anatomical and microscopic analysis that the cause of ITBS pain may be due to fat compression under the tract rather than friction against the lateral femoral epicondyle. The fact that these foundational pathology studies rank among the most cited highlights the community's reliance on robust anatomical data to inform management of patients with ITBS.

Our bibliometric analysis revealed that a majority of the papers included in our study were published relatively recently, with only 28% of papers being published before the year. The prevalence of more recent papers among the most cited articles is reflected in other studies on midshaft clavicle fractures (Tandron, Cohen, Cohen, et al. 2023) and acromioclavicular joint reconstruction (O'Dwyer et al. 2025). This finding may reflect both an advancement in research tools, growing clinical recognition of ITBS, and advances in 3D motion capture and musculoskeletal imaging. This may have allowed researchers to identify specific abnormalities in anatomy and biomechanics that caused the development of ITBS. These findings, assisted with newer developments of high-tech imaging and gait analysis, have allowed for the evolution of new gait-retraining protocols (Allen 2014) and treatment modalities (Sanchez-Alvarado et al. 2024).

This analysis also found a relative lack of papers associated with a higher LOE, with only three level I studies and two level II studies included in our analysis. This find-

ing was also reflected in bibliometric analyses of knee arthroscopy (Murphy et al. 2021) and femoral shaft fractures (Medlar et al. 2025). This gap in research could be explained by the heterogeneous presentation of ITBS patients, which may make it difficult to encompass all patients in inclusion criteria. In addition, long follow-up times for randomized controlled trials for newer treatment methods such as gait retraining may indicate that although there currently aren't many articles with higher LOE, there are randomized controlled trials possibly underway that are attempting to establish the long-term efficacy of newly established treatment methods. We predict that there may be an increase in articles published on ITBS that have an LOE of I or II.

There are a number of limitations in this study. Although the collection of data was cross-checked by two reviewers, there could still be bias and skew of data due to subjectivity of data collection. In addition, only Web of Science was searched; other highly-impactful articles on ITBS could be available for citation outside of this platform. Additionally, it is important to note that citation counts do not necessarily equate to quality or clinical impact. Lastly, educational materials like textbooks, abstracts and lectures were not included in this study, though they may be impactful and available for citation.

Submitted: August 11, 2025 EDT. Accepted: October 08, 2025 EDT. Published: May 21, 2026 EDT.



This is an open-access article distributed under the terms of the Creative Commons Attribution 4.0 International License (CCBY-NC-ND-4.0). View this license's legal deed at <https://creativecommons.org/licenses/by-nc-nd/4.0> and legal code at <https://creativecommons.org/licenses/by-nc-nd/4.0/legalcode> for more information.

REFERENCES

- Allen, D. J. 2014. "Treatment of Distal Iliotibial Band Syndrome in a Long Distance Runner with Gait Re-Training Emphasizing Step Rate Manipulation." *Int J Sports Phys Ther* 9 (2): 222–31.
- Beals, C., and D. Flanigan. 2013. "A Review of Treatments for Iliotibial Band Syndrome in the Athletic Population." *J Sports Med* 2013:1–6. <https://doi.org/10.1155/2013/367169>.
- Ellegaard, O. 2018. "The Application of Bibliometric Analysis: Disciplinary and User Aspects." *Scientometrics* 116 (1): 181–202. <https://doi.org/10.1007/s11192-018-2765-z>.
- Fairclough, J., K. Hayashi, H. Toumi, et al. 2006. "The Functional Anatomy of the Iliotibial Band during Flexion and Extension of the Knee: Implications for Understanding Iliotibial Band Syndrome." *J Anat* 208 (3): 309–16. <https://doi.org/10.1111/j.1469-7580.2006.00531.x>.
- Ferber, R., B. Noehren, J. Hamill, and I. Davis. 2010. "Competitive Female Runners With a History of Iliotibial Band Syndrome Demonstrate Atypical Hip and Knee Kinematics." *J Orthop Sports Phys Ther* 40 (2): 52–58. <https://doi.org/10.2519/jospt.2010.3028>.
- Fredericson, M., C. L. Cookingham, A. M. Chaudhari, B. C. Dowdell, N. Oestreich, and S. A. Sahrmann. 2000. "Hip Abductor Weakness in Distance Runners with Iliotibial Band Syndrome." *Clin J Sport Med* 10 (3): 169–75. <https://doi.org/10.1097/00042752-200007000-00004>.
- Marais, J. V., A. Jansen Van Rensburg, M. P. Schwellnus, E. Jordaan, and P. Boer. 2024. "Risk Factors Associated with a History of Iliotibial Band Syndrome (hITBS) in Distance Runners: A Cross-Sectional Study in 76 654 Race Entrants – a SAFER XXXIII Study." *Phys Sportsmed* 52 (6): 568–78. <https://doi.org/10.1080/00913847.2024.2341607>.
- Medlar, C., C. J. Kilkenny, M. S. Davey, F. Farooq, and B. J. O'Daly. 2025. "The Top 50 Cited Publications Relating to Femoral Shaft Fractures – A Bibliometric Analysis of the Literature." *J Orthop* 67:88–93. <https://doi.org/10.1016/j.jor.2025.01.025>.
- Murphy, S. N., M. L. Moore, J. R. Pollock, K. S. McQuivey, and J. S. Bingham. 2021. "The Top 50 Most-Cited Knee Arthroscopy Studies." *Arthrosc Sports Med Rehabil* 3 (4): e1243–53. <https://doi.org/10.1016/j.asmr.2021.05.006>.
- Noble, C. A. 1980. "Iliotibial Band Friction Syndrome in Runners." *Am J Sports Med* 8 (4): 232–34. <https://doi.org/10.1177/036354658000800403>.
- Noehren, B., I. Davis, and J. Hamill. 2007. "ASB Clinical Biomechanics Award Winner 2006." *Clin Biomech* 22 (9): 951–56. <https://doi.org/10.1016/j.clinbiomech.2007.07.001>.
- O'Dwyer, L., B. Murphy, M. S. Davey, D. Morrissey, and J. T. Cassidy. 2025. "A Bibliometric Analysis of the 50 Most Cited Articles on Acromioclavicular Joint Reconstruction." *Musculoskelet Surg*, January. <https://doi.org/10.1007/s12306-025-00886-w>.
- Pegrum, J., A. Self, and N. Hall. 2019. "Iliotibial Band Syndrome." *BMJ*, March, 1980. <https://doi.org/10.1136/bmj.1980>.
- Sanchez-Alvarado, A., C. Bokil, M. Cassel, and T. Engel. 2024. "Effects of Conservative Treatment Strategies for Iliotibial Band Syndrome on Pain and Function in Runners: A Systematic Review." *Front Sports Act Living* 6:1386456. <https://doi.org/10.3389/fspor.2024.1386456>.
- Tandron, M., L. Cohen, J. Cohen, et al. 2023. "The Fifty Most-Cited Articles Regarding Midshaft Clavicle Fractures." *J Orthop* 39:50–58. <https://doi.org/10.1016/j.jor.2023.03.014>.
- Wright, J., M. Swiontkowski, and J. Heckman. 2003. "Introducing Levels of Evidence to The Journal." *J Bone Jt Surg* 85 (1): 1–3. <https://doi.org/10.2106/00004623-200301000-00001>.